

# Exam 1 Formula Sheet

**Things you know but can't quite remember:**

$$\begin{aligned}
 g &= 9.81 \text{ m/s}^2 & \text{sphere } V &= \frac{4}{3}\pi r^3 & ax^2 + bx^2 + c = 0 \implies x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 \sin(\theta \pm \varphi) &= \sin \theta \cos \varphi \pm \cos \theta \sin \varphi & \cos(\theta \pm \varphi) &= \cos \theta \cos \varphi \mp \sin \theta \sin \varphi \\
 2 \cos \theta \cos \varphi &= \cos(\theta + \varphi) + \cos(\theta - \varphi) & 2 \sin \theta \sin \varphi &= \cos(\theta - \varphi) - \cos(\theta + \varphi) \\
 2 \sin \theta \cos \varphi &= \sin(\theta + \varphi) + \sin(\theta - \varphi) & c^2 &= a^2 + b^2 - 2ab \cos \theta_{ab} \\
 \cos \theta + \cos \varphi &= 2 \cos\left(\frac{\theta + \varphi}{2}\right) \cos\left(\frac{\theta - \varphi}{2}\right) & \cos \theta - \cos \varphi &= 2 \sin\left(\frac{\theta + \varphi}{2}\right) \sin\left(\frac{\theta - \varphi}{2}\right) \\
 \sin \theta + \sin \varphi &= 2 \sin\left(\frac{\theta + \varphi}{2}\right) \cos\left(\frac{\theta - \varphi}{2}\right) & \cos^2 \theta &= \frac{1}{2}(1 + \cos 2\theta) & \sin^2 \theta &= \frac{1}{2}(1 - \cos 2\theta) \\
 e^{i\theta} &= \cos \theta + i \sin \theta & \cos \theta &= \frac{1}{2}(e^{i\theta} + e^{-i\theta}) & \sin \theta &= \frac{1}{2}(e^{i\theta} - e^{-i\theta})
 \end{aligned}$$

**Those other ones:**

$$\cosh z = \frac{1}{2}(e^z + e^{-z}) = \cos iz \quad \sinh z = \frac{1}{2}(e^z - e^{-z}) = \sin iz \quad \tanh z = \frac{\sinh z}{\cosh z} \quad \cosh^2 z - \sinh^2 z = 1$$

**“What will this ever be good for?”**

$$\begin{aligned}
 \frac{d}{dx} \sin ax &= a \cos ax & \frac{d}{dx} \cos ax &= -a \sin ax & \frac{d}{dx} \sinh ax &= a \cosh ax & \frac{d}{dx} \cosh ax &= a \sinh ax \\
 \frac{d}{dx} \tan z &= \sec^2 z & \frac{d}{dx} \tanh z &= \operatorname{sech}^2 z & \frac{d}{dz} \sinh z &= \cosh z & \frac{d}{dz} \cosh z &= \sinh z \\
 \int u \, dv &= uv - \int v \, du & \int \frac{dx}{1+x^2} &= \arctan x & \int \frac{dx}{1-x^2} &= \operatorname{arctanh} x \\
 \int \frac{dx}{\sqrt{1-x^2}} &= \arcsin x & \int \frac{dx}{\sqrt{1+x^2}} &= \operatorname{arcsinh} x & \int \tan(x) \, dx &= -\ln \cos x & \int \tanh x \, dx &= \ln \cosh x \\
 \int \frac{dx}{x+x^2} &= \ln\left(\frac{x}{1+x}\right) & \int \frac{x \, dx}{1+x^2} &= \frac{1}{2} \ln(1+x^2) & \int \frac{dx}{\sqrt{x^2-1}} &= \operatorname{arccosh} x & \int \frac{x \, dx}{\sqrt{1+x^2}} &= \sqrt{1+x^2} \\
 \int \frac{dx}{x\sqrt{x^2-1}} &= \arccos(1/x) & \int \frac{\sqrt{x} \, dx}{\sqrt{1-x}} &= \arcsin(\sqrt{x}) - \sqrt{x(1-x)} & \int \frac{dx}{(1+x^2)^{3/2}} &= \frac{x}{(1+x^2)^{1/2}} \\
 \int \ln(x) \, dx &= x \ln(x) - x
 \end{aligned}$$

**Arrows:**

$$\begin{aligned}
 |\mathbf{F}| &= \sqrt{F_x^2 + F_y^2} & \text{magnitude} & \theta = \tan^{-1} \left[ \frac{F_y}{F_x} \right] & \text{direction} & \mathbf{a} \cdot \mathbf{b} = a_x b_x + a_y b_y + a_z b_z = |\mathbf{a}| |\mathbf{b}| \cos \theta \\
 \mathbf{a} \times \mathbf{b} &= \det \begin{bmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{bmatrix} = (a_y b_z - a_z b_y) \hat{\mathbf{x}} + (a_z b_x - a_x b_z) \hat{\mathbf{y}} + (a_x b_y - a_y b_x) \hat{\mathbf{z}}
 \end{aligned}$$

**Almost,  $x \ll 1$**

$$\begin{aligned}
 f(z) &= f(a) + f'(a)(z-a) + \frac{1}{2!} f''(a)(z-a)^2 + \frac{1}{3!} f'''(z-a)^3 + \dots \\
 (1+x)^n &\approx 1 + nx + \frac{1}{2} n(n+1)x^2 & \tan x &\approx x + \frac{1}{3}x^3 & \sin x &\approx x - \frac{1}{3!}x^3 & \cos x &\approx 1 - \frac{1}{2}x^2 \\
 e^x &\approx 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 & \ln(1+z) &\approx z - \frac{1}{2}z^2 & \text{should your calculator be in degrees or radians? ... just saying.}
 \end{aligned}$$

### Revenge of the Arrows:

$$\begin{aligned}\Sigma \mathbf{F} &= \mathbf{F}_{\text{net}} = m\mathbf{a} & \Sigma F_x &= ma_x & \Sigma F_y &= ma_y, \text{ etc.} & F_{\text{grav}} &= mg = \text{weight} & \mathbf{F}_{12} &= -\mathbf{F}_{21} \\ \mathbf{F} &= \frac{d\mathbf{p}}{dt} \approx \frac{\Delta(m\mathbf{v})}{\Delta t} & \text{direction } \dots & f_s \leq \mu_s n & f_k &= \mu_k n & \mathbf{F}_c &= -\frac{mv^2}{r} \hat{\mathbf{r}} \\ \mathbf{F} &= m\ddot{\mathbf{r}} = \begin{cases} F_x &= m\ddot{x} \\ F_y &= m\ddot{y} \text{ (Cartesian } x, y, z) \\ F_z &= m\ddot{z} \end{cases} = \begin{cases} F_r &= m(\ddot{r} - r\dot{\varphi}^2) \\ F_\varphi &= m(r\ddot{\varphi} + 2\dot{r}\dot{\varphi}) \end{cases} \text{ (2D polar } r, \varphi) \\ \mathbf{F}_{12} &= k_e \frac{q_1 q_2}{r_{12}^2} \hat{\mathbf{r}}_{12} = q_2 \mathbf{E}_1 & \mathbf{r}_{12} &= \mathbf{r}_1 - \mathbf{r}_2 & \mathbf{F}_B &= q\mathbf{v} \times \mathbf{B} & \mathbf{F} &= q(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \end{aligned}$$

### The Arrows Strike Back:

$$\begin{aligned}\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) &= \mathbf{B} \cdot (\mathbf{C} \times \mathbf{A}) = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B}) & \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) \\ \nabla f &= \hat{\mathbf{x}} \frac{\partial f}{\partial x} + \hat{\mathbf{y}} \frac{\partial f}{\partial y} + \hat{\mathbf{z}} \frac{\partial f}{\partial z} & \text{Cartesian} & \nabla f &= \hat{\mathbf{r}} \frac{\partial f}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial f}{\partial \theta} + \hat{\varphi} \frac{1}{r \sin \theta} \frac{\partial f}{\partial \varphi} & \text{Spherical} \\ \nabla \times \mathbf{F} &= \hat{\mathbf{x}} \left( \frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z} \right) + \hat{\mathbf{y}} \left( \frac{\partial F_x}{\partial z} - \frac{\partial F_z}{\partial x} \right) + \hat{\mathbf{z}} \left( \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) & \text{Cartesian} \\ \nabla \times \mathbf{F} &= \hat{\mathbf{r}} \frac{1}{r \sin \theta} \left[ \frac{\partial}{\partial \theta} (F_\varphi \sin \theta) - \frac{\partial F_\theta}{\partial \varphi} \right] + \hat{\theta} \left[ \frac{1}{r \sin \theta} \frac{\partial F_r}{\partial \varphi} - \frac{1}{r} \frac{\partial}{\partial r} (r F_\varphi) \right] + \hat{\varphi} \frac{1}{r} \left[ \frac{\partial}{\partial r} (r F_\theta) - \frac{\partial F_r}{\partial \theta} \right] & \text{Spherical} \end{aligned}$$

### Blowback:

$$f(v) = f_{\text{lin}} + f_{\text{quad}} \quad f_{\text{lin}} = bv \quad f_{\text{quad}} = cv^2 \quad \text{separate and integrate}$$

### Space Oddity:

$$\frac{dx}{dt} = \dot{x} \quad \frac{d^2x}{dt^2} = \frac{d\dot{x}}{dt} = \ddot{x} \quad m\dot{v} = -\dot{m}v_{\text{ex}} + F^{\text{ext}}$$

### Centering and Rotation:

$$\mathbf{R} = \frac{1}{M} \sum_{\alpha=1}^N m_\alpha \mathbf{r}_\alpha = \frac{m_1 \mathbf{r}_1 + \dots + m_N \mathbf{r}_N}{M} \quad \mathbf{l} = \mathbf{r} \times \mathbf{p} \quad \mathbf{L} = \sum_{\alpha=1}^N \mathbf{l}_\alpha = \sum_{\alpha=1}^N \mathbf{r}_\alpha \times \mathbf{p}_\alpha \quad \dot{\mathbf{L}} = \mathbf{F}^{\text{ext}}$$

### Work, meh:

$$\Delta T = T_2 - T_1 = \int_1^2 \mathbf{F} \cdot d\mathbf{r} \equiv W(1 \rightarrow 2) \quad U(\mathbf{r}) = -W(\mathbf{r}_o \rightarrow \mathbf{r}) \equiv - \int_{\mathbf{r}_o}^{\mathbf{r}} \mathbf{F}(\mathbf{r}') \cdot d\mathbf{r}' \quad \mathbf{F} = -\nabla U$$

central force:  $\mathbf{F} = f(\mathbf{r}) \hat{\mathbf{r}}$     central force is spherically symmetric [ $f(\mathbf{r})=f(r)$ ] if and only if conservative

### Interacting is difficult (conservative forces):

$$U = U^{\text{int}} + U^{\text{ext}} = \sum_{\alpha} \sum_{\beta > \alpha} U_{\alpha\beta} + \sum_{\alpha} U_{\alpha}^{\text{ext}} \quad (\text{net force on } \alpha) = -\nabla_{\alpha} U$$

### Wiggling:

$$F = -kx \iff U = \frac{1}{2} kx^2 \quad \ddot{x} = -\omega^2 x \iff x(t) = A \cos(\omega t - \delta)$$

$$\ddot{x} + 2\beta\dot{x} + \omega^2 x = 0 \iff x(t) = Ae^{-\beta t} \cos(\omega_1 t - \delta) \quad \beta = b/2m \quad \omega_o = \sqrt{\frac{k}{m}} \quad \omega_1 = \sqrt{\omega_o^2 - \beta^2} \quad (\beta < \omega_o)$$

$$\text{driven } A^2 = \frac{f_o^2}{(\omega_o - \omega)^2 + 4\beta^2 \omega^2} \quad \delta = \arctan \left( \frac{2\beta\omega}{\omega_o^2 - \omega^2} \right)$$

**Basic physics you should really know at this point, so I am printing it super small.**

Rotation: we use radians

1-D motion:

$$v(t) = \frac{d}{dt}x(t) \quad a(t) = \frac{d}{dt}v(t) = \frac{d^2}{dt^2}x(t)$$

$$v(t) = \int_0^t a \, dt \quad x(t) = \int_0^t v \, dt$$

↑ const. acc.

$$x_f = x_i + v_{xi}t + \frac{1}{2}a_x t^2$$

$$v_{xf}^2 = v_{xi}^2 + 2a_x \Delta x$$

$$v_f = v_i + at$$

Projectile motion:

$$v_x(t) = v_i \cos \theta \quad v_y(t) = v_i \sin \theta - gt$$

$$x(t) = x_i + v_x t \quad y(t) = y_i + v_y t + \frac{1}{2}a_y t^2$$

over level ground:

$$\text{max height } = H = \frac{v_i^2 \sin^2 \theta_i}{2g}$$

$$\text{Range } = R = \frac{v_i^2 \sin 2\theta_i}{g}$$

2-D motion:

$$\mathbf{r} = x(t)\hat{i} + y(t)\hat{j}$$

$$x(t) = x_i + v_{ix}t + \frac{1}{2}a_x t^2$$

$$y(t) = y_i + v_{iy}t + \frac{1}{2}a_y t^2$$

$$\mathbf{v} = v_x(t)\hat{i} + v_y(t)\hat{j}$$

$$v_x(t) = \frac{dx}{dt} = v_{xi} + a_x t$$

$$v_y(t) = \frac{dy}{dt} = v_{yi} + a_y t$$

$$\mathbf{a} = a_x(t)\hat{i} + a_y(t)\hat{j}$$

$$a_x(t) = \frac{dv_x}{dt} \quad a_y(t) = \frac{dv_y}{dt}$$

$$a_c = \frac{v^2}{r} \quad T = \frac{2\pi r}{v} \quad \text{circ.}$$

Force:

$$\Sigma \mathbf{F} = \mathbf{F}_{\text{net}} = m\mathbf{a} \quad \Sigma F_x = m a_x \quad \Sigma F_y = m a_y$$

$$F_{\text{grav}} = mg = \text{weight}$$

$$\mathbf{F}_{12} = -\mathbf{F}_{21}$$

$$\mathbf{F} = \frac{d\mathbf{p}}{dt} \approx \frac{\Delta(m\mathbf{v})}{\Delta t} \quad \text{direction ...}$$

$$f_s \leq \mu_s n \quad f_k = \mu_k n$$

$$\mathbf{F}_c = -\frac{mv^2}{r}\hat{r}$$

Work-Energy:

$$K = \frac{1}{2}mv^2 = \frac{p^2}{2m}$$

$$\Delta K = K_f - K_i = W$$

$$W = \int F(x) dx = -\Delta U \quad F = -\frac{dU}{dx} \quad \text{in equil.} \quad F = 0$$

$$U_g(y) = mgy \quad U_s(x) = \frac{1}{2}kx^2$$

$$K_i + U_i = K_f + U_f + W_{\text{ext}} = K_f + U_f + \int F_{\text{ext}} dx$$

$$s = \theta r \quad \leftarrow \text{arclength}$$

$$\omega = \frac{d\theta}{dt} = \frac{v}{r} \quad \alpha = \frac{d\omega}{dt} = \frac{d^2\theta}{dt^2}$$

$$a_t = \alpha r \quad \text{tangential} \quad a_r = \frac{v^2}{r} = \omega^2 r \quad \text{radial}$$

$$I = \sum_i m_i r_i^2 \Rightarrow \int r^2 dm = kmr^2$$

$$I_z = I_{\text{com}} + md^2 \quad \text{axis } z \text{ parallel, dist } d$$

$$\tau_{\text{net}} = \sum \tau = I\alpha = \frac{d\mathbf{L}}{dt}$$

$$\tau = \mathbf{r} \times \mathbf{F} \quad |\tau| = rF \sin \theta r_F$$

$$\mathbf{L} = \mathbf{r} \times \mathbf{p} = I\omega \quad L_i = L_f$$

$$K = \frac{1}{2}I\omega^2 = L^2/2I$$

$$\Delta K = \frac{1}{2}I\omega_f^2 - \frac{1}{2}I\omega_i^2 = W = \int \tau d\theta$$

$$P = \frac{dW}{dt} = \tau\omega$$

Gravitation:

$$\mathbf{F}_{12} = -\frac{Gm_1 m_2}{r^2} \hat{r}_{12} = -\frac{dU_g}{dr} \hat{r}$$

$$g = \frac{GM_e}{R_e^2}$$

$$U_g(r) = -\int F(r) dr = \frac{GMm}{r}$$

$$K + U_g \geq 0 \quad \text{escape} \quad K + U_g < 0 \quad \text{bound}$$

$$\frac{dA}{dt} = \frac{1}{2}r^2\omega = \frac{L}{2m} \quad T^2 = \left(\frac{4\pi^2}{GM}\right)r^3$$

$$E_{\text{orbit}} = \frac{-GMm}{2a} \quad \text{elliptical; } a \rightarrow r \text{ for circular}$$

Oscillations:

$$\omega = \frac{2\pi}{T} = 2\pi f \quad T = \frac{1}{f}$$

$$x(t) = x_m \cos(\omega t + \varphi)$$

$$a = \frac{d^2x}{dt^2} = -\omega^2 x \quad \text{for SHM}$$

$$\omega = \sqrt{k/m} \quad \text{spring}$$

$$U = -\frac{1}{2}kx^2 \quad F = -\frac{dU}{dx} = ma$$

$$T = \begin{cases} 2\pi\sqrt{I/\kappa} & \text{torsion pendulum} \\ 2\pi\sqrt{L/g} & \text{simple pendulum} \\ 2\pi\sqrt{T/mgd_{\text{cm}}} & \text{physical pendulum} \end{cases}$$