

Quiz 4: momentum and such

1. A rubber ball strikes a brick wall with a velocity (just before the collision) of $\vec{v}_i = 3.0\hat{i} + 4.0\hat{j}$ m/s. It rebounds with a velocity of $\vec{v}_f = -3.0\hat{i} + 4.0\hat{j}$ m/s, *i.e.*, the collision was perfectly elastic. What can be said about the change in momentum of the rubber ball?

- The momentum \vec{p} did not change, since the collision was elastic.
- The momentum \vec{p} changed direction, its magnitude is the same.
- The momentum \vec{p} decreased.
- The momentum \vec{p} increased.
- The question cannot be answered, we do not know the mass of the ball.

Solution: The magnitude of the momentum $|\vec{p}| = m|\vec{v}|$ is the same in either case, but its direction has changed. Both before and after the collision, $|\vec{v}| = 3^2 + 4^2 = 5^2$, so independent of the mass, $|\vec{p}|$ is constant.

2. Which of the following is in error?

- $\vec{F} = \frac{d\vec{p}}{dt}$
- $K = \frac{p^2}{2m}$
- $\vec{p} = m \frac{d\vec{x}}{dt}$
- $\Delta\vec{p} = \int \vec{F} \cdot d\vec{r}$

Solution: With $p = mv$ and constant mass m , all are true but the last option, it should be $\Delta\vec{p} = \int \vec{F} dt$.

3. A 60 kg person standing on a frictionless surface fires a 0.5 kg arrow horizontally at 50 m/s. With what velocity does the archer move backwards across the ice after firing the arrow?

- $v_{\text{archer}} = +0.42$ m/s
- $v_{\text{archer}} = -0.42$ m/s
- $v_{\text{archer}} = -0.84$ m/s
- $v_{\text{archer}} = +0.84$ m/s

Solution: Conservation of momentum: initially there is zero momentum, so afterwards we must have $m_{\text{archer}}v_{\text{archer}}+m_{\text{arrow}}v_{\text{arrow}} = 0$. The archer's velocity must be negative, ruling out two answers right off the bat. Solving for v_{archer} and plugging in the numbers, $v_{\text{archer}} = +0.42 \text{ m/s}$.

4. A ball of mass m is dropped from rest at a height h . What is the magnitude of the ball's momentum just before impact?

- $2mgh$
- $m\sqrt{gh}$
- $m\sqrt{2gh}$
- $\sqrt{2mgh}$

Solution: The ball starts at height h above the ground, with potential energy $U_i = mgh$. It reaches the ground with zero potential energy, and thus kinetic energy $K_f = \frac{1}{2}mv^2 = U_i = mgh$. This gives the velocity as the ball strikes the ground as $v = \sqrt{2gh}$, so the momentum is $p = mv = m\sqrt{2gh}$.